Brain Research, 510 (1990) 360–362
Elsevier

Author Index

Amassian, V.E., see Levy Jr., W.J., 130
Anderson, E.G., see Glaum, S.R., 12
Arase, K., see Shimizu, H., 216
Ault, B., see Caviedes, P., 229
Azmitia, E.C., Murphy, R.B. and Whitaker-Azmitia, P.M., MDMA (Ecstasy) effects on cultured serotonergic neurons: evidence Ca^{2+}-dependent toxicity linked to release, 97

Baba, S., see Inui, A., 211
Baganoff, M.P., see Watson, G.B., 158
Baker, H.J., see Walkley, S.U., 63
Barish, M.E. and Raissdana, S.S., Induction of class II major histocompatibility complex antigens on a population of astrocytes from a mouse strain (BALB/c) resistant to experimental allergic encephalomyelitis, 329
Beck, Th., Wree, A. and Schleicher, A., Postischemic glucose utilization in rat hippocampal layers, 74
Beltz, T.G., see Sullivan, M.J., 237
Bingmann, D., see Walden, J., 127
Blessing, W.W., see Halliday, G.M., 104
Blumbergs, P.C., see Halliday, G.M., 104
Bol, A., see Veraart, C., 115
Bolanowski, M.A., see Watson, G.B., 158
Bray, G.A., see Shimizu, H., 216
Bray, G.A., see Shimizu, H., 251
Burnard, D.M., Crichton, S.A. and MacVicar, B.A., Electrophysiological properties of reactive glial cells in the kainate-lesioned hippocampal slice, 43

Cadwell, J., see Levy Jr., W.J., 130
Calderón, F., see Torrealba, F., 351
Caviedes, P., Ault, B. and Rapoport, S.I., The role of altered sodium currents in action potential abnormalities of cultured dorsal root ganglion neurons from trisomy 21 (Down syndrome) human fetuses, 229
Changelian, P.S., Meiri, K., Soppet, D., Valenza, H., Loewy, A. and Willard, M., Purification of the growth-associated protein GAP-43 by reversed phase chromatography: amino acid sequence analysis and cDNA identification, 259
Chihara, K., see Inui, A, 211
Christie, W.M., see Maxwell, D.J., 346
Cohen, Y., see Robert, J.J., 7
Coleman, P.A., see Fohlmeister, J.F., 343
Consorti, D., see Fagioli, S., 150
Cook, J.E., Morphological recovery of axotomized goldfish retinal ganglion cells in an environment known to prevent retinotopic refinement of their regenerated tectal arbors, 181
Cooper, T.R., see Smith III, M.W., 242
Cotton, R.G.H., see Halliday, G.M., 104
Coyle, J.T., see Lowenstein, P.R., 122
Crichton, S.A., see Burnard, D.M., 43

Davrainville, J.L., see Khayari, A., 140
Del Bianco, E., see Geppetti, P., 58
Demierre, B., Martinou, J.-C. and Kato, A.C., Embryonic motoneurons grated into the adult CNS can differentiate and migrate, 355
Deppeler, C.L., see Watson, G.B., 158
De Volder, A.G., see Veraart, C., 115
Durie, E., see Sharif, N.A., 108

Fagioli, S., Consorti, D. and Pavone, F., Age-related cholinergic drug effects on analgesia in two inbred strains of mice, 150
Fohlmeister, J.F., Coleman, P.A. and Miller, R.F., Modeling the repetitive firing of retinal ganglion cells, 343
François, C., see Sadikot, A.F., 161

Geffen, L.B., see Halliday, G.M., 104
Geppetti, P., Del Bianco, E., Santicioli, P., Lippe, I.Th., Maggi, C.A. and Sicuteri, F., Release of sensory neuropeptides from dural venous sinuses of guinea pig, 58
Gerstein, G.L., see Gochin, P.M., 195
Gervais, F., see Xin, L., 144
Glaum, S.R., Proudfit, H.K. and Anderson, E.G., 5-HT_3 receptors modulate spinal nociceptive reflexes, 12
Gochin, P.M., Gerstein, G.L. and Kaltenbach, J.A., Dynamic temporal properties of effective connections in rat dorsal cochlear nucleus, 195
Goffinet, A.M., see Veraart, C., 115
Gotoh, M., see Iguchi, A., 321
Graziadei, P.P.C., see Takami, S., 339

Halliday, G.M., Blumbergs, P.C., Cotton, R.G.H., Blessing, W.W. and Geffen, L.B., Loss of brainstem serotonin- and substance P-containing neurons in Parkinson's disease, 104
Hara, H., Kato, H. and Kogure, K., Protective effect of α-tocopherol on ischemic neuronal damage in the gerbil hippocampus, 335
Hayaishi, O., see Uda, R.H., 26
Herrera, D.G. and Robertson, H.A., Application of potassium chloride to the brain surface induces the c-fos proto-oncogene: reversal by MK-801, 166
Hiroi, N. and White, N.M., The reserpine-sensitive dopamine pool mediates (+)-amphetamine-conditioned reward in the place preference paradigm, 33
Hong, J.S, see Stachowiak, M.K., 277
Horiguchi, S., see Uda, R.H., 26
Hubert, F., see Molotchnikoff, S., 223
Hyodo, M., see Uda, R.H., 26

Iguchi, A., Gotoh, M., Matsunaga, H., Yatomi, A., Uemura, K., Miura, H., Kunoh, Y., Tamagawa, T. and Sakamoto, N., Neither adrenergic nor cholinergic antagonists in the central nervous system affect 2-deoxy-D-glucose (2-DG)-induced hyperglycemia, 321
Inoue, T., see Inui, A., 211
Inui, A., Inoue, T., Nakajima, M., Okita, M., Sakatani, N., Okimura, Y., Chihara, K. and Baba, S., Brain neuropeptide Y in the control of adrenocorticotropic hormone secretion in the dog, 211
Ito, S., see Uda, R.H., 26

Jacquot, C., see Robert, J.J., 7
Joh, T.H., see Smith III, M.W., 242
Johnson, A.K., see Sullivan, M.J., 237
Jones, E.E. and Naftolin, F., Estrogen effects on the tuberoinfundibular dopaminergic system in the female rat brain, 84
Jope, R.S., see Ormandy, G.C., 53
Joyce, J.N., see Lowenstein, P.R., 122

Kaltenbach, J.A., see Gochin, P.M., 195
Kashiwayanagi, M., see Sugawara, M., 326
Kato, A.C., see Demierre, B., 355
Kato, H., see Hara, H., 335
Kato, N., see Tsunashima, K., 171
Katoh-Semba, R., see Sano, M., 269
Kawasaki, F., see Narita, K., 289
Khayari, A., Mesfioui, A., Math, F. and Davrainville, J.L., Na^+,K^+-ATPase in the rat olfactory bulb: evidence for higher enzymatic activity in the glomerular layer, 140
Kirino, T., Tamura, A. and Sano, K., Chronic maintenance of presynaptic terminals in gliotic hippocampus following ischemia, 17
Kirino, T., see Tamura, A., 154
Kita, H., see Narita, K., 289
Kitajima, S., see Sano, M., 269
Kitamura, R., see Yamamoto, T., 309
Kogure, K., see Hara, H., 335
Kunoh, Y., see Iguchi, A., 321
Kurihara, K., see Sugawara, M., 326

Lanthorn, T.H., see Watson, G.B., 158
Levy Jr., W.J., Amassian, V.E., Traad, M. and Cadwell, J., Focal magnetic coil stimulation reveals motor cortical system reorganized in humans after traumatic quadriplegia, 130
Li, Y.-P. and Lombardini, J.B., Guanidinoethanesulfonic acid — inhibitor of GABA uptake in rat cortical synaptosomes, 147
Lippe, I.Th., see Geppetti, P., 58
Loewy, A., see Changelian, P.S., 259
Lombardini, J.B., see Li, Y.-P., 147
Lowenstein, P.R., Joyce, J.N., Coyle, J.T. and Marshall, J.F., Striosomal organization of cholinergic and dopaminergic uptake sites and cholinergic M_1 receptors in the adult human striatum: a quantitative receptor autoradiographic study, 122

MacVicar, B.A., see Burnard, D.M., 43
Maggi, C.A., see Geppetti, P., 58
Marshall, J.F., see Lowenstein, P.R., 122

Martinou, J.-C., see Demierre, B., 355
Masui, A., see Tsunashima, K., 171
Masuo, Y., Pélaprat, D., Montagne, M.-N., Scherman, D. and Rostène, W., Regulation of neurotensin-containing neurons in the rat striatum and substantia nigra. Effects of unilateral nigral lesion with 6-hydroxydopamine on neurotensin content and its binding site density, 203
Math, F., see Khayari, A., 140
Matsunaga, H., see Iguchi, A, 321
Maxwell, D.J., Christie, W.M., Ottersen, O.P. and Storm-Mathisen, J., Terminals of group 1a primary afferent fibres in Clarke's column are enriched with L-glutamate-like immunoreactivity, 346
Meiri, K., see Changelian, P.S., 259
Mendell, J.R., see Sahenk, Z., 1
Mesfioui, A., see Khayari, A., 140
Michel, A.D., see Sharif, N.A., 108
Michel, Ch., see Veraart, C., 115
Miller, R.F., see Fohlmeister, J.F., 343
Miura, H., see Iguchi, A., 321
Molotchnikoff, S. and Hubert, F., Susceptibility of neurons in area 18a to blockade of area 17 in rats, 223
Montagne, M.-N., see Masuo, Y., 203
Mori, K., see Taoka, N., 190
Murphy, R.B., see Azmitia, E.C., 97

Naftolin, F., see Jones, E.E., 84
Nakajima, M., see Inui, A., 211
Narita, K., Kawasaki, F. and Kita, H., Mn and Mg influxes through Ca channels of motor nerve terminals are prevented by verapamil in frogs, 289
Nishikawa, T., Teramoto, T. and Shimizu, S., Effect of morphine on Na^+,K^+-ATPase from homogenate of synaptosomes and of synaptic membrane of rat cerebral cortex, 92

Oka, H., see Tamura, A., 154
Okimura, Y., see Inui, A., 211
Okita, M., see Inui, A., 211
Ormandy, G.C. and R.S. Jope, R.S., Inhibition of phosphoinositide hydrolysis by the novel neurotoxin β-*N*-oxalyl-L-α,β-diaminopropionic acid (L-BOAA), 53
Orosco, M., see Robert, J.J., 7
Ottersen, O.P., see Maxwell, D.J., 346

Parent, A., see Sadikot, A.F., 161
Pavone, F., see Fagioli, S., 150
Pélaprat, D., see Masuo, Y., 203
Proudfit, H.K., see Glaum, S.R., 12

Raissdana, S.S., see Barish, M.E., 329
Rapoport, S.I., see Caviedes, P., 229
Rattazzi, M.C., see Walkley, S.U., 63
Ray, P.E., Ruley, E.J. and Saavedra, J.M., Down-regulation of angiotensin II receptors in subfornical organ of young male rats by chronic dietary sodium depletion, 303
Rekling, J.C., Excitatory effects of thyrotropin-releasing hormone (TRH) in hypoglossal motoneurons, 179
Richardson, P.M., see Xin, L., 144
Robert, J.J., Orosco, M., Rouch, C., Cohen, Y. and Jacquot, C., Opposite dopaminergic activity in lateral and median hypothalamic nuclei in relation to the feeding effect of D-Ser^2-Leu-Enk-Thr^6 (DSLET), 7
Robertson, H.A., see Herrera, D.G, 166
Rostène, W., see Masuo, Y., 203
Rouch, C., see Robert, J.J., 7
Ruley, E.J., see Ray, P.E., 303

Saavedra, J.M., see Ray, P.E., 303
Sadikot, A.F., Parent, A. and François, C., The centre médian and parafascicular thalamic nuclei project respectively to the sensorimotor and associative–limbic striatal territories in the squirrel monkey, 161
Sahenk, Z., Whitaker, J.N. and Mendell, J.R., Immunocytochemical evidence for the retrograde transport of intraaxonal cathepsin D: possible relevance to the dying-back process, 1
Sakamoto, N., see Iguchi, A, 321
Sakatani, N., see Inui, A., 211
Sano, K., see Kirino, T., 17
Sano, K., see Tamura, A., 154
Sano, M., Katoh-Semba, R., Kitajima, S. and Sato, C., Changes in levels of microtubule-associated proteins in relation to the outgrowth of neurites from PC12D cells, a forskolin- and nerve growth-responsive subline of PC12 pheochromocytoma cells, 269
Santicioli, P., see Geppetti, P., 58
Sato, C., see Sano, M., 269
Scherman, D., see Masuo, Y., 203
Schleicher, A., see Beck, Th., 74
Shargill, N.S., see Shimizu, H., 216
Sharif, N.A., Durie, E., Michel, A.D. and Whiting, R.L., Dog cerebral cortex contains μ-, δ-and κ-opioid receptors at different densities: apparent lack of evidence for subtypes of the κ-receptor using selective radioligands, 108
Shimizu, H. and Bray, G.A., Effects of insulin on hypothalamic monoamine metabolism, 251
Shimizu, H., Shargill, N.S., Arase, K., York, D.A. and Bray, G.A., Relationship between uptake of norepinephrine by hypothalamic homogenates and the activity of brown adipose tissue, 216
Shimizu, S., see Nishikawa, T., 92
Shimo-oku, M., see Taoka, N., 190
Sicuteri, F., see Geppetti, P., 58
Skamene, E., see Xin, L., 144
Smith, D.E., see Smith III, M.W., 242
Smith III, M.W., Cooper, T.R., Joh, T.H. and Smith, D.E., Cell loss and class distribution of TH-I cells in the substantia nigra of the neurological mutant, weaver, 242
Soppet, D., see Changelian, P.S., 259
Speckmann, E.-J., see Walden, J., 127
Stachowiak, M.K., Hong, J.S. and Viveros, O.H., Coordinate and differential regulation of phenylethanolamine *N*-methyltransferase, tyrosine hydroxylase and proenkephalin mRNAs by neural and hormonal mechanisms in cultured bovine adrenal medullary cells, 277
Starke, K., see Wichmann, T., 296
Storm-Mathisen, J., see Maxwell, D.J., 346
Straub, H., see Walden, J., 127
Sugawara, M., Kashiwayanagi, M. and Kurihara, K., Water response of frog olfactory system is induced by a decrease in osmotic pressure, 326
Sullivan, M.J., Beltz, T.G. and Johnson, A.K., Amastatin potentiation of drinking induced by blood-borne angiotensin: evidence for mediation by endogenous brain angiotensin, 237

Takagi, K., see Tamura, A., 154
Takami, S. and Graziadei, P.P.C., Morphological complexity of the glomerulus in the rat accessory olfactory bulb — a Golgi study, 339
Tamagawa, T., see Iguchi, A., 321
Tamura, A., Kirino, T., Sano, K., Takagi, K. and Oka, H., Atrophy of the ipsilateral substantia nigra following middle cerebral artery occlusion in the rat, 154
Tamura, A., see Kirino, T., 17
Taoka, N., Mori, K., Yamagata, Y. and Shimo-oku, M., Cervical afferent pathway to inferior oblique motoneuron in the cat, 190
Teramoto, T., see Nishikawa, T., 92
Torrealba, F. and Calderón, F., Central projections of coarse and fine vagal axons of the cat, 351
Traad, M., see Levy Jr., W.J., 130
Tsunashima, K., Masui, A. and Kato, N., The effect of delta sleep-inducing peptide (DSIP) and phosphorylated DSIP (P-DSIP) on the apomorphine-induced hypothermia in rats, 171

Uda, R.H., Horiguchi, S., Ito, S., Hyodo, M. and Hayaishi, O., Nociceptive effects induced by intrathecal administration of prostaglandin D_2, E_2, or $F_{2\alpha}$ to conscious mice, 26
Uemura, K., see Iguchi, A., 321

Valenza, H., see Changelian, P.S., 259
Veraart, C., De Volder, A.G., Wanet-Defalque, M.C., Bol, A., Michel, Ch. and Goffinet, A.M., Glucose utilization in human visual cortex is abnormally elevated in blindness of early onset but decreased in blindness of late onset, 115
Viveros, O.H., see Stachowiak, M.K., 277

Walden, J., Speckmann, E.-J., Bingmann, D. and Straub, H., Augmentation of *N*-methyl-D-aspartate induced depolarizations by GABA in neocortical and archicortical neurons, 127
Walkley, S.U., Wurzelmann, S., Rattazzi, M.C. and Baker, H.J., Distribution of ectopic neurite growth and other geometrical distortions of CNS neurons in feline GM2 gangliosidosis, 63
Wanet-Defalque, M.C., see Veraart, C., 115
Watson, G.B., Bolanowski, M.A., Baganoff, M.P., Deppeler, C.L. and Lanthorn, T.H., D-Cycloserine acts as a partial agonist at the glycine modulatory site of the NMDA receptor expressed in *Xenopus* oocytes, 158

Weerasuriya, A., Patterns of change in endoneurial capillary permeability and vascular space during nerve regeneration, 135
Whitaker, J.N., see Sahenk, Z., 1
Whitaker-Azmitia, P.M., see Azmitia, E.C., 97
White, N.M., see Hiroi, N., 33
Whiting, R.L., see Sharif, N.A., 108
Wichmann, T. and Starke, K., Modulation by muscarine and opioid receptors of acetylcholine release in slices from striatostriatal grafts in the rat, 296
Willard, M., see Changelian, P.S., 259
Wree, A., see Beck, Th., 74
Wurzelmann, S., see Walkley, S.U., 63

Xin, L., Richardson, P.M., Gervais, F. and Skamene, E., A deficiency of axonal regeneration in C57BL/6J mice, 144

Yamagata, Y., see Taoka, N., 190
Yamamoto, T. and Kitamura, R., A search for the cortical gustatory area in the hamster, 309
Yatomi, A., see Iguchi, A, 321
York, D.A., see Shimizu, H., 216

(contents continued)

Modeling the repetitive firing of retinal ganglion cells
J.F. Fohlmeister, P.A. Coleman and R.F. Miller (U.S.A.) 343
Terminals of group 1a primary afferent fibres in Clarke's column are enriched with L-glutamate-like immunoreactivity
D.J. Maxwell, W.M. Christie, O.P. Ottersen and J. Storm-Mathisen (U.K., Norway) 346
Central projections of coarse and fine vagal axons of the cat
F. Torrealba and F. Calderón (Chile) 351
Embryonic motoneurons grated into the adult CNS can differentiate and migrate
B. Demierre, J.-C. Martinou and A.C. Kato (Switzerland) 355

Author Index 360